

Chemotaxis to Furan Compounds by Furan-Degrading Pseudomonas Strains

Nancy N. Nichols, Tristan A. Lunde, Kevin C. Graden, Kate
A. Hallock, Cara K. Kowalchyk, Rebecca M. Southern, Ellen
J. Soskin and Jayna L. Ditty

Appl. Environ. Microbiol. 2012, 78(17):6365. DOI:
10.1128/AEM.01104-12.

Published Ahead of Print 22 June 2012.

Updated information and services can be found at:
<http://aem.asm.org/content/78/17/6365>

These include:

REFERENCES

This article cites 37 articles, 18 of which can be accessed free
at: <http://aem.asm.org/content/78/17/6365#ref-list-1>

CONTENT ALERTS

Receive: RSS Feeds, eTOCs, free email alerts (when new
articles cite this article), [more»](#)

Information about commercial reprint orders: <http://journals.asm.org/site/misc/reprints.xhtml>
To subscribe to to another ASM Journal go to: <http://journals.asm.org/site/subscriptions/>

Chemotaxis to Furan Compounds by Furan-Degrading *Pseudomonas* Strains

Nancy N. Nichols,^a Tristan A. Lunde,^b Kevin C. Graden,^b Kate A. Hallock,^b Cara K. Kowalchuk,^b Rebecca M. Southern,^b Ellen J. Soskin,^b and Jayna L. Ditty^b

Bioenergy Research Unit, USDA–National Center for Agricultural Utilization Research, Peoria, Illinois, USA,^a and Department of Biology, University of St. Thomas, St. Paul, Minnesota, USA^b

Two *Pseudomonas* strains known to utilize furan derivatives were shown to respond chemotactically to furfural, 5-hydroxymethylfurfural, furfuryl alcohol, and 2-furoic acid. In addition, a LysR-family regulatory protein known to regulate furan metabolic genes was found to be involved in regulating the chemotactic response.

Furan derivatives are heterocyclic, five-membered aromatic rings, consisting of four carbon atoms and one oxygen, that are used industrially as solvents and chemical precursors (5). Although naturally occurring in the environment (35), they are also formed by the Maillard reaction, and for this reason furan aldehydes are present in biomass sugar preparations that have been pretreated with dilute acid. This pretreatment method, which involves incubation of fibrous biomass in dilute H₂SO₄ under heat and pressure, generates furfural and 5-hydroxymethylfurfural (5-HMF) from pentose and hexose sugars originating from hemicellulose and cellulose (12). Although furan aldehydes are microbial inhibitors, a number of microbes can utilize these compounds as a source of carbon and energy (2–4, 11, 19, 33–37, 39). The ability of selected microbes to dissimilate furfural and 5-HMF is the basis for a biological abatement strategy for detoxification of microbial inhibitors, including furfural and 5-HMF, in biomass hydrolysates (19, 22, 24, 37).

An aerobic pathway for furfuryl alcohol (FA), furfural, and 2-furoic acid (2-FA) degradation was initially proposed for *Pseudomonas putida* Fu1 and *Pseudomonas* F2 (13, 14, 34), and some accessory genes involved in furan metabolism were cloned from *P. putida* Fu1 (23). The complete complement of genes for conversion of furfural and 5-HMF to 2-oxoglutarate has been cloned from *Cupriavidus basilensis* HMF14 (15). The pathway includes enzymes for the sequential oxidation of FA and furfural to 2-FA, along with a furoyl-coenzyme A synthetase and a molybdenum-dependent dehydrogenase. Following lactone hydrolysis, which may occur spontaneously, removal of the coenzyme A moiety is catalyzed by 2-oxoglutaroyl coenzyme A thioesterase. Metabolism of 5-HMF also proceeds via furoic acid (15).

Motile bacteria are capable of chemotaxis to a variety of carbon sources. Paradigms were established initially for chemotaxis to sugars and amino acids by *Escherichia coli*, *Salmonella*, and *Bacillus subtilis* (30), with an increasing diversity of chemotactic mechanisms having been described more recently (16, 21). The broad chemotactic capability of pseudomonads, long noted for metabolic diversity (31), has also been recognized (28). Chemotaxis of *Pseudomonas* has been reported for aromatic acids (8), chlorinated benzoates (7), nitro- and aminobenzoates (26), benzene, toluene, and trichloroethylene (10, 27), naphthalene (6), atrazine (17), 2,4-dichlorophenoxyacetate (2,4-D) (9), and pyrimidines (18), and the importance of chemotaxis in bioremediation efforts has recently been addressed (25, 29, 32). Chemotaxis to furan

compounds has not been reported previously; here, we describe the behavioral response of two *Pseudomonas* strains to furfural, 5-HMF, FA, and 2-FA.

Pseudomonas putida Fu1 (13) and *Pseudomonas* sp. strain A3 (19), both originally isolated for their ability to degrade furan molecules, were inoculated into 0.3% soft-agar minimal medium swarm plates (28) containing 0.5 mM 5-HMF, 1.0 mM furfural, or 0.5 mM 2-FA and incubated at 30°C for 24 to 36 h. As the cells metabolize the low concentration of attractant present in the soft agar, they follow a self-generated concentration gradient, resulting in the presence of a sharp ring of growth that spreads from the original point of inoculation and indicating a chemotactic response. Swarm plate assays (Fig. 1) demonstrated that strains Fu1 and A3 are attracted to furan molecules or potentially their metabolic intermediates, as metabolism is required for this assay. Subsequently, agarose plug assays (28, 38) were used to assess the ability of these strains to detect furans in a temporal manner over 10 minutes. Overnight cultures of *Pseudomonas* sp. strain A3 and *P. putida* Fu1 were grown in 5 mM 2-FA, harvested, and resuspended in chemotaxis buffer (CB) (7) to an A₆₆₀ of 1.0. Motile cell suspensions were added to a chamber such that they surrounded a low-melting-temperature agarose plug containing CB (negative control), 2% Casamino Acids (CA; positive control), or 1.0 mM 2-FA or FA. As the attractant diffuses out from the agarose plug into the surrounding environment, and if the cells are attracted to the compound, cells accumulate at the optimal attractant concentration, resulting in a characteristic white ring of cells that typically forms around the periphery of the agarose plug. Each white cell ring was assessed microscopically to ensure that the cells that had accumulated were motile. Initial images taken at time zero for both strains and all attractants showed the same diffuse suspension of cells. After a 10-min incubation at room temperature, a white ring of motile cells developed around the agarose plugs containing CA and 2-FA, while areas closest to the plug were devoid of cells. This result demonstrates that the cells can detect 2-FA in a

Received 5 April 2012 Accepted 14 June 2012

Published ahead of print 22 June 2012

Address correspondence to Jayna L. Ditty, jlditty@stthomas.edu.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/AEM.01104-12




FIG 1 *Pseudomonas putida* strain Fu1 and *Pseudomonas* sp. strain A3 are attracted to furan molecules. Soft-agar minimal medium swarm plates containing 0.5 mM 5-HMF, 1.0 mM furfural, or 0.5 mM 2-FA were inoculated with cells at the center of the swarm plate and incubated at 30°C for 24 to 36 h. As cells metabolize the attractant, they follow a self-generated concentration gradient, forming white concentric circles of cellular growth. Succinate (1.0 mM) swarm plates served as positive controls. Images were taken using back-lighting to visualize the outward growth patterns generated by attracted cells. Images shown are representative of assays conducted in triplicate.

spatial gradient and are not reliant on a self-generated gradient based on growth on 2-FA (Fig. 2A). *P. putida* Fu1 was also shown to detect FA under the same assay conditions (Fig. 2B).

The inducible nature of chemotactic responses to aromatic hydrocarbons and acids has been well established (29); therefore, quantitative capillary assays (1, 8) were conducted to determine if the response to 2-FA in *P. putida* Fu1 was inducible. Cells grown in




FIG 2 *Pseudomonas putida* strain Fu1 and *Pseudomonas* sp. strain A3 grown on 2-FA respond to a spatial gradient of furan molecules. (A). Strains Fu1 and A3 were grown on 5.0 mM 2-FA to provide inducing conditions. Cultures grown overnight with aeration were harvested and resuspended in chemotaxis buffer (CB) at an A_{600} of 1.0. Low-melting-point agarose (2%, with Coomassie brilliant blue for contrast) contained CB (negative control), 10% Casamino Acids (CA; positive control), or 1.0 mM 2-FA. A characteristic white ring of cells that develops at the optimal attractant concentration diffusing from the agarose plug demonstrates chemotactic behavior and is indicated in these images by a white arrow. Images were documented after 10 min by photography using back-lighting. Images shown are representative of assays conducted in triplicate. (B) *P. putida* strain Fu1 is attracted to furfuryl alcohol (FA) under the same assay conditions.




FIG 3 The response of *Pseudomonas putida* Fu1 to 2-FA is inducible. Aerated cultures were grown in 10 mM pyruvate (light gray bars) or 10 mM 2-FA (dark gray bars) at 30°C for 21 h. Cells were harvested, resuspended in CB, and placed in a U-tube chamber constructed by placing a U-shaped glass tube between a slide and a coverslip. Cells were allowed to chemotactically respond to either CB-, 2% CA-, or 2-FA-filled capillaries for 30 min at 30°C. Error bars represent the standard deviations for at least four independent assays conducted in triplicate. Means with the same letter are not significantly different ($P < 0.05$; one-way analysis of variance [ANOVA] interaction, Tukey multiple comparison test).

10 mM pyruvate (PYR) or 2-FA were harvested, washed, and resuspended in CB to an A_{600} of 1.0. The motile cell suspension was inserted into a chamber containing 1- μ l capillary tubes loaded with CB, 2% CA, or 10-fold dilutions of 2-FA ranging from 500 to 0.05 mM. The cells were allowed to chemotactically respond to the attractant diffusing from the capillaries for 30 min at 30°C. The capillary tubes were then removed from the chambers, and the contents were emptied in order to quantitate (by serial dilution and growth on agar plates) the number of bacteria that had responded by swimming toward the attractant and into the capillary. As shown in Fig. 3, the response of *P. putida* Fu1 to 2-FA is inducible by growth on 2-FA, as the number of cells that responded to 2-FA at any concentration when grown in the presence of PYR was not significantly different from the number responding to buffer alone. However, *P. putida* Fu1 chemotactically responded to an optimum concentration of 5 mM 2-FA when grown on 2-FA.

As shown in Fig. 2, the responses of strains Fu1 and A3 were qualitatively different, even when the strains were responding to a positive control such as CA. It has been our experience that different strains can respond differently to the same chemoattractants, as some strains are naturally more motile than others. This was evident in this study, as strain A3 clearly had a weaker overall chemotactic response to both CA and 2-FA in qualitative assays (Fig. 1 and 2). Strain A3 was therefore not tested further in the quantitative capillary assays, because of its overall weaker response.

Because the chemotactic response of *P. putida* Fu1 to 2-FA was shown to be inducible, we investigated the role in chemotaxis of a regulatory protein, PsfB, known to affect the expression of genes involved in furan metabolism (23). Strain PSF2 is a kanamycin-resistant mutant derived from Fu1 by minitransposon mutagenesis. PSF2 harbors the transposon in *psfB*, a LysR-type regulatory gene, and is incapable of growth on furfural and 2-FA (23). LysR




FIG 4 A LysR-family regulatory protein known to be responsible for the regulation of 2-FA metabolic genes regulates the inducible chemotactic response to 2-FA. *P. putida* strains Fu1, PSF2 (*psfB*::Km-inactivated strain), and PSF2 pPSF210 (*psfB*-complementing strain) were grown on 10 mM pyruvate (PYR) and 5.0 mM 2-FA to provide inducing conditions and assessed by the plug assay (see the legend to Fig. 2 for a detailed assay description). Attractants are listed to the left of each row. Descriptions of the *psfB* mutant PSF2 and the complementing plasmid pPSF210, which carries *psfB* in *trans*, can be found in reference 23. Images were documented after 10 min by photography using backlighting. Images shown are representative of assays conducted in triplicate. White arrows indicate positive chemotaxis responses.

family regulators typically activate transcription in the presence of an inducing molecule (20, 23). When grown in inducing conditions (PYR plus 2-FA), strain PSF2 did not respond to 2-FA, as assessed by agarose plug assays (Fig. 4.) However, when *PsfB* was produced in *trans* in mutant PSF2 under inducing conditions, the chemotactic response was restored to wild-type levels. As shown in Fig. 4A, a slight positive chemotactic response to 2-FA was also detected when the complemented strain was grown under noninducing conditions (absence of 2-FA). This positive response is consistent with the fact that *PsfB* is homologous to LysR-type positive regulators.

The results of this work clearly demonstrate that *Pseudomonas* strains capable of metabolizing furan molecules also have the capacity to detect and respond chemotactically to furans in their environment. Knowledge of the chemotactic response to furans explains the initial facet of bacterial degradation of these industrially important and naturally occurring compounds. The discovery of bacterial chemotaxis to furfural, 2-FA, FA, and 5-HMF expands the known chemotactic repertoire of pseudomonads and is proof of a behavioral response to a class of compounds, furans, that had not previously been examined. The future identification of microbial furan chemoreceptors will provide a more thorough understanding of the furan chemosensory system and allow comparison to the diversity of chemotaxis pathways and chemoreceptors identified *in silico* (21).

ACKNOWLEDGMENTS

We are grateful to Jan R. Andreesen for providing *P. putida* Fu1. The strain is also available as DSM6384 from the German Collection of Microorganisms and Cell Cultures (www.dsmz.de). We thank Lois Fruen, director of the Breck School Advanced Science Research Program, Minneapolis, MN, for supporting the work of C.K.K., R.M.S., and E.J.S.

This study was supported in part by a grant to J.L.D. from the National Science Foundation (MCB-0919930).

Mention of trade names or commercial products in this article is solely for the purpose of providing scientific information and does not imply recommendation or endorsement by the U.S. Department of Agriculture.

REFERENCES

- Adler J. 1973. A method for measuring chemotaxis and use of the method to determine optimum conditions for chemotaxis by *Escherichia coli*. J. Gen. Microbiol. 74:77–91.
- Boopathy R, Bokang H, Daniels L. 1993. Biotransformation of furfural and 5-hydroxymethylfurfural by enteric bacteria. J. Ind. Microbiol. Biotechnol. 11:147–150.
- Brune G, Schoberth SM, Sahn H. 1983. Growth of a strictly anaerobic bacterium on furfural (2-furaldehyde). Appl. Environ. Microbiol. 46:1187–1192.
- Folkerts M, et al. 1989. *Desulfovibrio furfuralis* sp. nov., a furfural degrading strictly anaerobic bacterium. Syst. Appl. Microbiol. 11:161–169.
- Gandini A, Belgacem MN. 1997. Furans in polymer chemistry. Prog. Polym. Sci. 22:1203–1379.
- Grimm AC, Harwood CS. 1997. Chemotaxis of *Pseudomonas* spp. to the polycyclic aromatic hydrocarbon naphthalene. Appl. Environ. Microbiol. 63:4111–4115.
- Harwood CS, Parales RE, Dispensa M. 1990. Chemotaxis of *Pseudomonas putida* toward chlorinated benzoates. Appl. Environ. Microbiol. 56:1501–1503.
- Harwood CS, Rivelli M, Ornston LN. 1984. Aromatic acids are chemoattractants for *Pseudomonas putida*. J. Bacteriol. 160:622–628.
- Hawkins AC, Harwood CS. 2002. Chemotaxis of *Ralstonia eutropha* JMP134(pJP4) to the herbicide 2,4-dichlorophenoxyacetate. Appl. Environ. Microbiol. 68:968–972.
- Kim HE, et al. 2006. Identification and characterization of the chemotactic transducer in *Pseudomonas aeruginosa* PAO1 for positive chemotaxis to trichloroethylene. J. Bacteriol. 188:6700–6702.
- Kim T, Hah Y, Hong S. 1983. Toxic effects of furfural on *Pseudomonas fluorescens*. Korean J. Microbiol. 21:149–155.
- Klinke HB, Thomsen AB, Ahring BK. 2004. Inhibition of ethanol-producing yeast and bacteria by degradation products produced during pre-treatment of biomass. Appl. Microbiol. Biotechnol. 66:10–26.
- Koenig K, Andreesen JR. 1989. Molybdenum involvement in aerobic degradation of 2-furoic acid by *Pseudomonas putida* Fu1. Appl. Environ. Microbiol. 55:1829–1834.
- Koenig K, Andreesen JR. 1990. Xanthine dehydrogenase and 2-furoyl-coenzyme A dehydrogenase from *Pseudomonas putida* Fu1: two molybdenum-containing dehydrogenases of novel structural composition. J. Bacteriol. 172:5999–6009.
- Koopman F, Wierckx N, De Winde JH, Ruijsenaars HJ. 2010. Identification and characterization of the furfural and 5-(hydroxymethyl)furfural degradation pathways of *Cupriavidus basilensis* HMF14. Proc. Natl. Acad. Sci. U. S. A. 107:4919–4924.
- Krell T, et al. 2011. Diversity at its best: bacterial taxis. Environ. Microbiol. 13:1115–1124.
- Liu X, Parales RE. 2009. Bacterial chemotaxis to atrazine and related s-triazines. Appl. Environ. Microbiol. 75:5481–5488.
- Liu X, Wood PL, Parales JV, Parales RE. 2009. Chemotaxis to pyrimidines and identification of a cytosine chemoreceptor in *Pseudomonas putida*. J. Bacteriol. 191:2909–2916.
- López MJ, Nichols NN, Dien BS, Moreno J, Bothast RJ. 2004. Isolation of microorganisms for biological detoxification of lignocellulosic hydrolysates. Appl. Microbiol. Biotechnol. 64:125–131.
- Maddocks SE, Oyston PC. 2008. Structure and function of the LysR-type transcriptional regulator (LTTR) family proteins. Microbiology 154:3609–3623.
- Miller LD, Russell MH, Alexandre G. 2009. Diversity in bacterial chemotactic responses and niche adaptation. Adv. Appl. Microbiol. 66:53–75.
- Nichols NN, Dien BS, Cotta MA. 2010. Fermentation of bioenergy crops into ethanol using biological abatement for removal of inhibitors. Bioreour. Technol. 101:7545–7550.
- Nichols NN, Mertens JA. 2008. Identification and transcriptional profiling of *Pseudomonas putida* genes involved in furoic acid metabolism. FEMS Microbiol. Lett. 284:52–57.
- Nichols NN, et al. 2008. Fungal metabolism of fermentation inhibitors present in corn stover dilute acid hydrolysate. Enzyme Microb. Technol. 42:624–630.

25. Pandey G, Jain RK. 2002. Bacterial chemotaxis toward environmental pollutants: role in bioremediation. *Appl. Environ. Microbiol.* **68**:5789–5795.
26. Parales RE. 2004. Nitrobenzoates and aminobenzoates are chemoattractants for *Pseudomonas* strains. *Appl. Environ. Microbiol.* **70**:285–292.
27. Parales RE, Ditty JL, Harwood CS. 2000. Toluene-degrading bacteria are chemotactic towards the environmental pollutants benzene, toluene, and trichloroethylene. *Appl. Environ. Microbiol.* **66**:4098–4104.
28. Parales RE, Harwood CS. 2002. Bacterial chemotaxis to pollutants and plant-derived aromatic molecules. *Curr. Opin. Microbiol.* **5**:266–273.
29. Parales RE, Ju K-S, Rollefson JB, Ditty JL. 2008. Bioavailability, transport, and chemotaxis towards organic pollutants, p 146–187. In Diaz E (ed), *Microbial biodegradation: genomics and molecular biology*. Caister Academic Press, Norfolk, United Kingdom.
30. Rao CV, Kirby JR, Arkin AP. 2004. Design and diversity in bacterial chemotaxis: a comparative study in *Escherichia coli* and *Bacillus subtilis*. *PLoS Biol.* **2**:E49. doi:10.1371/journal.pbio.0020049.
31. Stanier RY, Palleroni NJ, Doudoroff M. 1966. The aerobic pseudomonads: a taxonomic study. *J. Gen. Microbiol.* **43**:159–271.
32. Strobel KL, et al. 2011. Chemotaxis increases vertical migration and apparent transverse dispersion of bacteria in a bench-scale microcosm. *Biotechnol. Bioeng.* **108**:2070–2077.
33. Thiemer B, Andreesen JR, Schröder T. 2003. Cloning and characterization of a gene cluster involved in tetrahydrofuran degradation in *Pseudonocardia* sp. strain K1. *Arch. Microbiol.* **179**:266–277.
34. Trudgill PW. 1969. The metabolism of 2-furoic acid by *Pseudomonas* F2. *Biochem. J.* **113**:577–587.
35. Trudgill PW. 1984. The microbial metabolism of furans, p 295–308. In Gibson DT (ed), *Microbial degradation of organic compounds*. Marcel Dekker, Inc., New York, NY.
36. Wang P, Brenchley JE, Humphrey AE. 2004. Screening microorganisms for utilization of furfural and possible intermediates in its degradative pathway. *Biotechnol. Lett.* **16**:977–982.
37. Wierckx N, Koopman F, Bandounas L, De Winde JH, Ruijsenaars HJ. 2010. Isolation and characterization of *Cupriavidus basilensis* HMF14 for biological removal of inhibitors from lignocellulosic hydrolysate. *Microb. Biotechnol.* **3**:336–343.
38. Yu HS, Alam M. 1997. An agarose-in-plug bridge method to study chemotaxis in the archaeon *Halobacterium salinarum*. *FEMS Microbiol. Lett.* **156**:265–269.
39. Zarnt G, Schröder T, Andreesen JR. 1997. Degradation of tetrahydrofurfuryl alcohol by *Ralstonia eutropha* is initiated by an inducible pyrroloquinoline quinone-dependent alcohol dehydrogenase. *Appl. Environ. Microbiol.* **63**:4891–4898.